

# CLASS III DENTAL CARIES AND NEED FOR ROOT CANAL TREATMENT IN MAXILLARY ANTERIORS - A HOSPITAL BASED RETROSPECTIVE ANALYSIS

<sup>1</sup>S.S.Shivanni, <sup>2</sup>Dr. Anjaneyulu K\*, <sup>3</sup>Dr. Balakrishna R.N

<sup>1</sup>Saveetha Dental College and Hospitals, Saveetha Institute Of Medical and Technical Science, Saveetha University, Chennai -600 077.

<sup>2</sup>\*Reader, Department of Conservative Dentistry and Endodontics, Saveetha Dental College and Hospitals, Saveetha Institute Of Medical and Technical Science, Saveetha University, Chennai-600 077.

<sup>3</sup>Saveetha Dental College and Hospitals, Saveetha Institute Of Medical and Technical Science, Saveetha University, Chennai-600077

DOI: 10.47750/pnr.2022.13.S04.162

The aim of this study is to assess the number of RCT done in the maxillary anteriors among patients with Class III caries. The study includes patients visiting the OP of Saveetha Dental College diagnosed with class III caries on the maxillary anteriors and to find the number of RCTs done among those patients. The results were statistically analysed using SPSS software. Descriptive statistics and chi-square tests were done for gender and diagnosis of the tooth. Among a total of 345 patients diagnosed with class III caries on the maxillary anteriors, 21.74% have undergone RCT while 78.26% of the patients did not undergo RCT. The incidence of RCT done in patients with Class III caries was found to be low. This might be due to the aesthetic consideration of the patients and hence early treatment.

**KEYWORDS:** Irreversible pulpitis, RCT, class III caries, maxillary anteriors.

## INTRODUCTION:

Dental caries is one of the most common preventable diseases, people are susceptible to the disease throughout their lifetime.(Pitts, 2004)(Fejerskov and Kidd, 2003)(Pitts, 2004) It can be arrested and can be reversed in its early stages, but is often not self limiting and without proper care, caries can progress until the tooth is lost. (Sciences and US Department of Health and Human Services; National Institutes of Health; National Institute on General Medical Sciences, 2000)-(Kidd, Giedrys-Leeper and Simons, 2000)

Dental caries is the localised destruction of susceptible dental hard tissues by acidic by-products from bacterial fermentation of carbohydrates.(Fejerskov and Kidd, 2003)(Marsh and Martin, 1992)(Fejerskov and Kidd, 2003) The decayed surface is the sequela of the disease process and is a sign of advanced disease. (Kidd, Thylstrup and Fejerskov, 1981) The signs of the carious demineralisation are seen on the hard dental tissues, but the disease process is initiated within the bacterial biofilm that covers a tooth surface. Moreover, the very early changes in the enamel are not detected with traditional clinical and radiographic methods. Dental caries is a multifactorial disease, several factors play a role in the initiation and progression of the lesion including, environmental, host and behavioral factors.(Marsh and Martin, 1992; Shaffer *et al.*, 2013)

To know how early in the caries process the pulp may be involved is important for the pathogenesis of dental caries and treatment. So far no systemic examination has been made of the response of the pulp in early caries. The effect of dental caries on the pulp, impairment of odontoblasts and inflammation of the pulp were observed only under very deep active lesions in the dentin. No observations of pulpal response under enamel caries alone was made.(Brännström and Lind, 1965)

Root canal therapy (RCT) and tooth extraction have been conventional treatment options for management of human mature teeth with irreversible pulpitis.(Asgary *et al.*, 2013) Root canal treatment is a common procedure in dentistry. The main indications for RCT are irreversible pulpitis and necrosis of dental pulp due to caries or dental trauma.(R. Rajakeerthi and Ms, 2019)

Commented [1]: Change title in DOC name

Now the growing trend in this area motivated us to pursue this project. Various aspects of endodontics and conservative dentistry over the past five years. Now we are focusing on retrospective studies, the idea for which has stemmed from the current interest in our community. Our team has extensive knowledge and research experience that has translate into high quality publications (Sathivel *et al.*, 2008; Panda *et al.*, 2014; Govindaraju, Neelakantan and Gutmann, 2017; Johnson *et al.*, 2020; Saraswathi *et al.*, 2020) (Kumar *et al.*, 2006; Devi and Gnanavel, 2014; Varghese *et al.*, 2015; Sivamurthy and Sundari, 2016; Chen *et al.*, 2019). This study aims in finding the prevalence of class III caries among maxillary anteriors and the incidence of those anteriors with an infected pulp that underwent root canal therapy.

## MATERIALS AND METHODS :

### Study population:

A retrospective study was carried out among 345 patients who visited University in Chennai diagnosed with class III caries on the maxillary anteriors. Among these, the prevalence of root canal treatment done was assessed. The data was collected from patients records between June 2019-March 2020. The data contains details of patients, intraoral photographs and treatment being done.

### Inclusion criteria:

- Patients with class III caries without pulpal involvement
- Patients with class III caries who underwent RCT
- Maxillary anteriors

### Exclusion criteria:

- Patients with periradicular disease
- Pulp necrosis

**Sample size:** Sample size [N=345] is the total number of patients who visit University in Chennai with class III caries. Their distribution according to gender and diagnosis were recorded.

**Ethical approval:** Ethical clearance was obtained from the Institutional Ethical Committee and Scientific Review Board [SRB] of University in Chennai.

**Data analysis:** The data collected were entered in an Excel sheet and subjected to statistical analysis using SPSS software. Descriptive statistics was done i.e frequency and cross tabulation. A chi square test was done between gender and diagnosis of the tooth. Independent variables are gender while dependent variables are diagnosis of the tooth. The level of significance was  $p < 0.05$ .

## RESULTS AND DISCUSSION:

It was found that male distribution was predominant with 65.80% of cases whereas females were only about 34.20% [Figure 1]. The prevalence of class III caries in maxillary anteriors without pulpal involvement was found to be 78.26%, whereas maxillary anteriors with class III caries that had pulpal involvement and had undergone RCT was only 21.74% of the cases [Figure 2]. Figure 3 shows the association between gender and diagnosis among the population. It was found that Class III caries in maxillary anteriors that have not undergone RCT was highly prevalent among males (55.36%) than females (22.90%), whereas Class III caries in maxillary anteriors that have undergone RCT was found to be higher in females (11.30%) than males (10.43%). chi-square test shows high significance ( $p$  value = 0.000).

The present study was conducted to evaluate the prevalence of maxillary anteriors with class III caries with pulp involvement among both genders. The results of this study shows that there was a higher percentage of class III caries without pulp involvement [78.26%] and lower incidence of RCT performed due to class III caries [21.74%] in maxillary anteriors. There is no previous literature that shows the association between class III caries and Root canal therapy. However, the prevalence of class III caries have been studied. A study by Mustang Demirci *et al.*, shows that the prevalence of proximal caries in incisors, canines, premolars and occlusal fissures in molars was the highest in both sexes. (Demirci, Tuncer and Yuceokur, 2010) A study by Talabani RM *et al.*, shows that 331 [4.1 %] of patients had class III mesial caries and 289 [3.6%] of patients had class III distal caries. (Talabani, Al-Zahawi and Ibrahim, 2015) A similar study done in Nepal state that the prevalence of class III caries among 18.75 %. (Bhagat and Shrestha, 2014)

The limitation of this study is that it is done among a limited population and geographic location. The future scope of the study is to conduct the study among a larger population and to consider the prognosis. Our institution is passionate about high quality evidence based research and has excelled in various fields (Pc, Marimuthu and Devadoss, 2018; Ramesh *et al.*, 2018; Ezhilarasan, Apoorva and Ashok Vardhan, 2019; Ramadurai *et al.*, 2019;

Sridharan *et al.*, 2019; Vijayashree Priyadharsini, 2019; Mathew *et al.*, 2020). We hope this study adds to this rich legacy.

**Figures**




Figure 1: Shows the gender distribution among the study population. X axis corresponds to the gender and Y axis corresponds to the number of patients with class III dental caries. Male (green) distribution was higher when compared to females (purple) with a percentage of 65.80% and 34.20% respectively.




Figure 2: Bar chart shows the comparison between number of patients with normal class III dental caries and number of patients with Class III dental caries that had undergone RCT. X axis corresponds to diagnosis of patients with class III caries and Y axis corresponds to the number of patients. Number of patients with class III caries which have not undergone RCT (blue) were higher with a percentage of 78.26% when compared to the number of patients with class III caries which have undergone RCT (red) with a percentage of 21.74%.




Figure 3: Bar chart shows the association between gender and diagnosis. X axis corresponds to the gender and Y axis corresponds to the number of patients. Number of patients with class III caries that had not undergone RCT (blue) was more in males (55.36%) when compared to females (22.90%), whereas number of patients with class III caries that had undergone RCT (red) was found to be higher in females (11.30%) when compared to males (10.43%). The difference was statistically significant (chi-square value - 13.488, p value - 0.000 (< 0.05)).

### CONCLUSION :

The current study shows highest prevalence of maxillary anteriors with class III caries without pulp involvement and that underwent Root canal therapy. This may be because patients might consider treatment of class III caries in maxillary anteriors in the earlier stages as the aesthetics is compromised. Among the patients with class III caries who had undergone RCT, females were found to be significantly higher than males.

**AUTHOR CONTRIBUTIONS:** All authors have equal contributions in bringing out this research work.

**ACKNOWLEDGEMENT:** This study was supported by Saveetha Dental College and Hospitals, Chennai.

**CONFLICT OF INTEREST:** Nil.

### REFERENCES:

1. Asgary, S. et al. (2013) 'One-year results of vital pulp therapy in permanent molars with irreversible pulpitis: an ongoing multicenter, randomized, non-inferiority clinical trial', *Clinical Oral Investigations*, pp. 431–439. doi:10.1007/s00784-012-0712-6.
2. Bhagat, T.K. and Shrestha, A. (2014) 'Prevalence of dental caries among public school children in the Eastern Nepal', *Journal of Chitwan Medical College*, pp. 30–32. doi:10.3126/jcmc.v4i1.10845.
3. Brännström, M. and Lind, P.O. (1965) 'Pulpal Response to Early Dental Caries', *Journal of Dental Research*, pp. 1045–1050. doi:10.1177/00220345650440050701.
4. Chen, F. et al. (2019) '6-shogaol, a active constituents of ginger prevents UVB radiation mediated inflammation and oxidative stress through modulating NF2 signaling in human epidermal keratinocytes (HaCaT cells)', *Journal of photochemistry and photobiology. B, Biology*, 197, p. 111518.
5. Demirci, M., Tuncer, S. and Yuceokur, A.A. (2010) 'Prevalence of Caries on Individual Tooth Surfaces and its Distribution by Age and Gender in University Clinic Patients', *European Journal of Dentistry*, pp. 270–279. doi:10.1055/s-0039-1697839.
6. Devi, V.S. and Gnanavel, B.K. (2014) 'Properties of Concrete Manufactured Using Steel Slag', *Procedia Engineering*, 97, pp. 95–104.
7. Ezhilarasan, D., Apoorva, V.S. and Ashok Vardhan, N. (2019) 'Syzygium cumini extract induced reactive oxygen species-mediated apoptosis in human oral squamous carcinoma cells', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(2), pp. 115–121.
8. Fejerskov, O. and Kidd, E. (2003) *Dental Caries: The Disease and Its Clinical Management*. Wiley-Blackwell.
9. Govindaraju, L., Neelakantan, P. and Gutmann, J.L. (2017) 'Effect of root canal irrigating solutions on the compressive strength of tricalcium silicate cements', *Clinical oral investigations*, 21(2), pp. 567–571.
10. Johnson, J. et al. (2020) 'Computational identification of MiRNA-7110 from pulmonary arterial hypertension (PAH) ESTs: a new microRNA that links diabetes and PAH', *Hypertension research: official journal of the Japanese Society of Hypertension*, 43(4), pp. 360–362.
11. Kidd, E.A.M., Giedrys-Leeper, E. and Simons, D. (2000) 'Take Two Dentists: A Tale of Root Caries', *Dental Update*, pp. 222–230. doi:10.12968/denu.2000.27.5.222.
12. Kidd, E.A.M., Thylstrup, A. and Fejerskov, O. (1981) 'The Histopathology of Enamel Caries in Fluorosed Deciduous Teeth', *Caries Research*, pp. 346–352. doi:10.1159/000260537.
13. Kumar, M.S. et al. (2006) 'Expression of matrix metalloproteinases (MMP-8 and -9) in chronic periodontitis patients with and without diabetes mellitus', *Journal of periodontology*, 77(11), pp. 1803–1808.
14. Marsh, P. and Martin, M. (1992) 'Oral Microbiology'. doi:10.1007/978-1-4615-7556-6.
15. Mathew, M.G. et al. (2020) 'Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary ...', *Clinical oral investigations* [Preprint]. Available at: <https://link.springer.com/article/10.1007/s00784-020-03204-9>.
16. Panda, S. et al. (2014) 'Platelet rich fibrin and xenograft in treatment of intrabony defect', *Contemporary clinical dentistry*, 5(4), pp. 550–554.
17. Pe, J., Marimuthu, T. and Devadoss, P. (2018) 'Prevalence and measurement of anterior loop of the mandibular canal using CBCT: A cross sectional study', *Clinical implant dentistry and related research* [Preprint]. Available at: <https://europepmc.org/article/med/29624863>.
18. Pitts, N.B. (2004) 'Are We Ready to Move from Operative to Non-Operative/Preventive Treatment of Dental Caries in Clinical Practice?', *Caries Research*, pp. 294–304. doi:10.1159/000077769.
19. Ramadurai, N. et al. (2019) 'Effectiveness of 2% Articaine as an anesthetic agent in children: randomized controlled trial', *Clinical oral*

Commented [2]: Conclusion in accordance of aim

- investigations, 23(9), pp. 3543–3550.
20. Ramesh, A. *et al.* (2018) 'Comparative estimation of sulfiredoxin levels between chronic periodontitis and healthy patients - A case-control study', *Journal of periodontology*, 89(10), pp. 1241–1248.
  21. R, R., Rajakeerthi, R. and Ms, N. (2019) 'Natural Product as the Storage medium for an avulsed tooth – A Systematic Review', *Cumhuriyet Dental Journal*, pp. 249–256. doi:10.7126/cumudj.525182.
  22. Saraswathi, I. *et al.* (2020) 'Impact of COVID-19 outbreak on the mental health status of undergraduate medical students in a COVID-19 treating medical college: a prospective longitudinal study', *PeerJ*, p. e10164. doi:10.7717/peerj.10164.
  23. Sathivel, A. *et al.* (2008) 'Anti-peroxidative and anti-hyperlipidemic nature of Ulva lactuca crude polysaccharide on D-galactosamine induced hepatitis in rats', *Food and chemical toxicology: an international journal published for the British Industrial Biological Research Association*, 46(10), pp. 3262–3267.
  24. Sciences, U.S.D. of H.A.H.S.N.I. of H.N.I. on G.M. and US Department of Health and Human Services; National Institutes of Health; National Institute on General Medical Sciences (2000) 'The Chemistry of Health', *PsycEXTRA Dataset* [Preprint]. doi:10.1037/e400662005-001.
  25. Shaffer, J.R. *et al.* (2013) 'GWAS of Dental Caries Patterns in the Permanent Dentition', *Journal of Dental Research*, pp. 38–44. doi:10.1177/0022034512463579.
  26. Sivamurthy, G. and Sundari, S. (2016) 'Stress distribution patterns at mini-implant site during retraction and intrusion—a three-dimensional finite element study', *Progress in orthodontics*, 17(1), pp. 1–11.
  27. Sridharan, G. *et al.* (2019) 'Evaluation of salivary metabolomics in oral leukoplakia and oral squamous cell carcinoma', *Journal of oral pathology & medicine: official publication of the International Association of Oral Pathologists and the American Academy of Oral Pathology*, 48(4), pp. 299–306.
  28. Talabani, R.M., Al-Zahawi, A. and Ibrahim, R.O. (2015) 'Prevalence And Distribution Of Dental Caries Experience According To GV Black Classification For Patient Attending To Dental School', *Journal of Oral Health and Community Dentistry*, pp. 60–63. doi:10.5005/johcd-9-2-60.
  29. Varghese, S.S. *et al.* (2015) 'Estimation of salivary tumor necrosis factor-alpha in chronic and aggressive periodontitis patients', *Contemporary clinical dentistry*, 6(Suppl 1), pp. S152–6.
  30. Vijayashree Priyadharsini, J. (2019) 'In silico validation of the non-antibiotic drugs acetaminophen and ibuprofen as antibacterial agents against red complex pathogens', *Journal of periodontology*, 90(12), pp. 1441–1448.