

Effects of Covid 19 on Health status and Quality of life among Software Professionals in a Home based work setting - A Questionnaire based Cross Sectional Study

Dr. J. Vineeta Debbie Nesam¹, Dr. S. Saradha², P. Indhra³

¹Assistant Professor, Department of Pharmacology, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education, Kelambakkam, Chengalpattu district, Tamil Nadu, India.

²Professor, Department of Pharmacology, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education, Kelambakkam, Chengalpattu district, Tamil Nadu, India.

³Biostatistician, Department of Community Medicine, Chettinad Hospital and Research Institute, Chettinad Academy of Research and Education, Kelambakkam, Chengalpattu district, Tamil Nadu, India.

Email: saradhachoks@gmail.com

DOI: 10.47750/pnr.2023.14.03.065

Abstract

Introduction: Coronavirus disease (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), was first identified in December 2019 in Wuhan city, China. Following the outbreak of COVID-19 pandemic, to restrict the spread of virus, the Government has imposed strict measures with subsequent lockdowns as a result the home environment has become a workplace for many. Of all sectors the software domain is the most affected. Limited resource facility, work - engagement, work-life balance, family conflicts, stress management have been the challenges faced by these IT professionals working from home

Aim: To analyze the impact of Covid 19 on health status and quality of life among software professionals in Covid pandemic through validated online questionnaires using google forms.

Materials and Methods: This cross-sectional e-survey was conducted using validated questionnaires through google forms following approval of the Institutional Ethics Committee. The questionnaire consisted of 4 open ended and 21 close ended questions. Job-stress, physical and mental well-being, work life balance, work engagement was assessed. The google form was posted in social media channel groups of software professionals for a period of 2 weeks following which the data was collected. Statistical analysis was done using SPSS software version 22.0.

Results: A total of 154 software professionals participated in the study. Of which 80 were females and belonged to the 26-30 years age group. 83% software professionals preferred working in hybrid mode.

This study shows that working hours increased with increased work pressure in a home-based work setting (61%). Both work life balance and work engagement were fair in a home-based work setting (p-0.0001). Anxiety, depression and emotional problems affected work engagement and work life balance. Some professionals developed physical ailments such as hair fall, vision problems and back ache in a home-based work set up.

Conclusion: The findings of this study suggests that emotional well-being, sleep quality, decision latitude, limited resource facilities and job stress affected work engagement and work life balance. Hence undertaking appropriate measures proactively to enhance these factors under crisis conditions may mitigate the negative consequences of home-based work setting and improve the quality of life of software developers in the covid pandemic.

Keywords: Job stress, physical, mental well-being, work engagement and work life balance.

INTRODUCTION

Coronavirus disease (COVID-19) caused by Severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2), was first identified in Wuhan city, China in December 2019. On January 30, 2020, COVID-19, was declared a public health emergency situation by the WHO. Consequently, it became a pandemic that affected millions of people all over the world.^[1] There are 54.1 crore cases and 63.2 lakh Covid deaths worldwide. The first SARS-CoV-2 positive case in India was reported in the state of Kerala on January 30, 2020. Currently there are 83990 active cases and 524941 deaths in India.

Following the outbreak of COVID-19 pandemic the Government has imposed strict measures to restrict the spread of virus

with subsequent lockdowns which has shut all the industries, factories, IT companies, schools, colleges, tourism, transportation between states, countries and this resulted in the huge loss to government as well the private sectors with fall in economy.

With the measures undertaken during the pandemic, the home environment has become a workplace for parents, a school for children, and a living space for the whole family at the same time. Of all sectors the software domain is most affected as these IT professionals are subjected to work for long hours with limited resource facility in a home based work setting and software development needs high level of concentration, focused attention, being systematic, persistent, creative and need to withstand stress. This poses a challenge especially when deadlines are demanding to finish the assigned projection time affecting work-life balance, physical and psychological well-being and productivity of software professionals.^[2]

This forced work from home owing to Covid pandemic has aggravated occupational health problems like musculoskeletal disorders, vision problems, headache, psycho-social and sleep disorders. Work - engagement, work-life balance, limited resource facility, family conflicts, stress management have been the challenges faced by these IT professionals working from home.^[4] Moreover, this pandemic will have long-lasting effects in the software industry, allowing their employees to work from home for an indefinite period. Therefore, it is crucial to analyze how a home-based work setting affects software developers' well-being and productivity. This study will also provide insight on measures to be taken to mitigate the negative effects of work from home on mental being and productivity of software professionals.

Aim:

To analyze the impact of Covid 19 on health status and quality of life among software professionals in a Home-based work setting during Covid pandemic.

Objectives:

1. Primary Objective: To evaluate the effects of Covid 19 on physical and mental well-being of Software Professionals in a Home-based work setting.
2. Secondary Objective: To evaluate the quality of life of Software Professionals in a Home-based work setting.

Inclusion criteria:

1. Software professionals of the IT sector working from home were included in the study.
2. Age 22-40 years of either sex.
3. Participants willing to participate in the study.

Exclusion Criteria:

1. Age > 45 years.
2. Other professions working from home.
3. Software professionals of higher rank such as Directors, Vice Presidents, Chief Executive Officers, Chief Information Officers were excluded.
4. Software professionals with co-morbid illness.
5. Those with Psychiatric Illness on chronic treatment.

MATERIALS AND METHODS:

This cross-sectional study was conducted to study the impact of Covid 19 pandemic on health status among software professionals working from home through validated online questionnaires using Google forms following approval of the Institutional Human Ethics Committee. The questionnaire was validated by experts of 4 different fields. Informed consent was obtained from the participants prior commencement of the study. To assess the impact of Covid 19 on health status and quality of life of software professionals working from home a questionnaire was developed with 4 open ended questions and 21 close ended questions. Job stress, physical, mental well-being, work life balance, work engagement were the domains assessed. The Google forms were posted in social media channel groups of software professionals for a period of 2 weeks following which the data was collected and statistically analyzed using SPSS software version 22.

Statistical Analysis:

The data was analyzed using SPSS version 22. Descriptive statistics was done for all data and were expressed as percentages.

Categorical data was analyzed using Pearson's Chi square to find the correlation between the dependent variables. P value<0.05 was considered as significant.

RESULTS:

A total of 154 software professionals participated in the study. Of which 80 were female [Table:2]. Most of them belonged to the 26-30 years age group [Figure:1]. 63% have been working in a home based setting for the past 2 years [Table:4]. High frequency of software professionals (83) preferred working in hybrid mode which involved working a few days at home and a few days at office in a week. [Table:3].

Figure 1: Age-wise distribution




Table 2: Sex wise Distribution

Sex	Frequency	Percent
Female	80	51.9
Male	74	48.1
Total	154	100.0

Table 3: Working Preferences

Working Preferences	Frequency	Percentage
Hybrid mode	83	53.9
Work in home based setting	51	33.1
Working in office set up	20	13
Total	154	100

Table 4: Duration of work in home-based setting

	Frequency	Percent
< 1 year	16	10.4
> 2 year	97	63.0
1-2 year	41	26.6
Total	154	100.0

This study shows that 59% Software professionals worked for more hours (9-12 hours) a day in a home-based work setting compared to working in an office set up (6-9 hours) [Figure:6] with increased work pressure at home (61%) [Table:5] and 51% software professionals felt the need of quitting the job due to overall increased stress [Table:7]. They also felt the interaction with superiors was better working in an office set up (88%) but felt the ease of completing a task satisfactorily in a home based work setting (81%) [Table:8].

Table 5: Increased Work pressure during work in home-based setting

	Frequency	Percent
No	59	38.3
Yes	95	61.7
Total	154	100.0

Figure 6: Working hours during work in home-based setting and in office set up




Table 7: Feel like quitting job due to the stress of work in home-based setting

	Frequency	Percent	Valid Percent	Cumulative Percent
No	74	48.1	48.1	48.1
Yes	80	51.9	51.9	100.0
Total	154	100.0	100.0	

Table 8: Task assigned completed satisfactorily

Work mode	Task assigned completed satisfactorily		Total
	Working from Home	Working in office set up	
Hybrid mode	41	42	83
Working from Home	38	13	51
Working in office set up	2	18	20
Total	81	73	154

Chi-square=24.718^a, p=0.0001

65% software professionals denied receiving any increment during the covid imposed forced work from home. 80% of them suffered changes in sleep pattern. Sleeping hours has increased in a home-based work setting with p value 0.0001 [Figure:9].




Figure 9: Sleeping hours during work in home-based setting and in office set up

Time spent in physical fitness activities was less than one hour in both work from home and in the office set up [Table:10]. An opportunity for leisure activities was better in a home-based work setting compared to office set up. Screen time including laptop (P value-0.0001), Television and mobile (P value -0.02) was comparatively increased in a home-based work setting [Table:11,12]. Some professionals developed physical ailments of which most commonly encountered was hair fall, vision problems and back ache in a home-based work set up. They also complained of increased weight gain in a home-based work setting (73%).

Time spent on Physical Fitness during work in Home based setting	Time spent on Physical Fitness during work in office set up		Total
	< 1 hours	1-2 hours	
< 1 hours	69	38	107
> 2 hours	4	2	6
1-2 hours	30	11	41
Chi-Square 1.010, p -0.604			
Table 10: Time spent on Physical Fitness during work in Home-based setting and Office set up			

Using Laptop during work in Home-based setting	Using Laptop during work at Office set up				Total
	< 4 hours	>12 hours	4-8 hours	8-12 hours	
< 4 hours	4	0	4	0	8
>12 hours	0	2	4	5	11
4-8 hours	1	1	24	14	40
8-12 hours	7	1	55	32	95
Pearson Chi-Square-36.150, p-.0001					
Table 11: Using Laptop during work in Home-based setting and at Office set up					

Mobile usage during work in Home-based setting and at Office set up	Using Mobile at Office setup			Total
	< 2 hours	> 4 hours	2-4 hours	
<4 hours	64	6	21	91
>12 hours	4	0	2	6
4-8 hours	24	3	17	44
8-12 hours	3	3	7	13
Total	95	12	47	154
Pearson Chi-Square 14.153 ^a p-.028				
Table 12: Mobile usage during work in Home-based setting and at Office set up				

Both work life balance and work engagement which was fair in 54 software professionals with P value 0.0001 [Fig/Table:13]. 61% software professionals had anxiety and depression while working in a home-based setting which led to a fair work life

balance (P value 0.0001) [Fig/Table:14]. Emotional problems also interfered with work life balance and most of them could not fulfill their family responsibilities. (P value 0.0001). [Table:15].

Work-life balance	Work engagement				Total
	Excellent	Fair	Good	Poor	
Excellent	2	3	5	0	10
Fair	2	54	17	5	78
Good	4	12	23	3	42
Poor	0	12	5	7	24
Chi-Square -40.300 ^a , p-.0001					
Table 13: Correlation between work life balance and work engagement during work in home based setting					

Work-life balance	Anxious and depressed				Total
	All of the time	Most of the time	None of the time	Some of the time	
Excellent	0	0	5	5	10
Fair	3	18	2	55	78
Good	4	8	7	23	42
Poor	3	10	0	11	24
Total	10	36	14	94	154
Chi-Square 38.955 ^a , p-.0001					
Table 14: Effect of Anxiety & depression on work life balance					

Work-life balance	Emotional problems interfering with your Work in a home based work setting				Total
	All of the time	Most of the time	None of the time	Some of the time	
Excellent	0	1	7	2	10
Fair	0	21	5	52	78
Good	2	12	6	22	42
Poor	1	12	2	9	24
Total	3	46	20	85	154
Chi-Square-42.463, p-.0001					
Table 15: Emotional problems interfering with Work in a home based work setting					

DISCUSSION:

154 software professionals participated in this study of which 80 were females of 26-30 years of age. As shown in table 5 the overall stress was increased working from home due to inability to focus owing to interruptions from family members at home affecting work life balance which was similar to the study done by Denae Ford et al. (2021).

Working from home has also increased the working hours with reduced sleep hours leading to increased job stress and has increased the level of anxiety, depression and physical ailments affecting the well-being of these software professionals. It has also decreased the time spent in physical fitness and leisure activities similar to the study done by D.Ford et al. (2020)

While working remotely at home people tend to over work neglecting their daily routines, family responsibilities, no time for leisure activities and physical fitness leading to imbalance in work life balance and leading to poor quality of life and productivity at work. Hence by adopting new strategies like scheduling meetings, developing new daily routines, engaging in hobbies, spending time with family and friends we can improve the work environment and quality of life of software professionals working in a home set up as mentioned by Daniel Russo et al in his study. (2021)

Improving the quality of sleep which is an important predictor of well-being, reducing the workload by software companies and by providing increments and promotions as mentioned in a similar study done by Paul Ralf et al, (2020) we can improve the productivity and emotional well-being of the software professionals working from home.

Adopting stress reduction measures like meditation, yoga, slow breathing exercises and social interactions can reduce stress levels as mentioned by Naik et al, (2018) we can improve the work performance and quality of life of these software professionals. Thus, by improving these predictors of well-being and productivity by adopting appropriate measures we can remove the negative consequences of work from home and improve the quality of life of these software professionals in the near future.

CONCLUSION:

The findings of this study suggests that emotional well-being, physical ailments, sleep quality, decision latitude, limited resource facilities, inadequate increments and job stress affect work engagement and work life balance substantially. Hence undertaking appropriate measures proactively to enhance these factors under crisis conditions may mitigate the negative consequences of home-based work setting on software developers and provide benefits in terms of mental well-being and work engagement so as to improve the quality of life of software professionals as similar pandemic periods are foreseen in the future and home has become an alternative work set up for software developers and many software organizations.

ACKNOWLEDGEMENTS:

All authors are thankful to Chettinad Hospital and Research Institute and Chettinad Academy of Research and Education for facilitating the conduct of this research. We immensely thank all IT professionals who participated in the survey.

CONFLICT OF INTEREST: We have no conflicts of interest to disclose.

AUTHOR FUNDING: No external funding.

REFERENCES:

1. Daniel Russo, Paul H. P. Hanel: Predictors of well-being and productivity among software professionals during the COVID-19 pandemic – a longitudinal study: Empirical Software Engineering, Empirical Software Engineering, Volume 26, Article number:2021, volume 26, Article number: 62.
2. Gul Tokdemir: Software professionals during the COVID-19 pandemic in Turkey: Factors affecting their mental well-being and work engagement in the home-based work setting: Journal of Systems and Software: June 2022, Volume 188, 111286.
3. Ralph et al. "Pandemic programming: How COVID-19 affects software developers and how their organizations can help" Empirical Software Engineering 2020.
4. D. Ford M.-A. Storey T. Zimmermann C. Bird S. Jaffe C. Maddila et al. "A tale of two cities: Software developers working from home during the COVID-19 pandemic" arXiv preprint arXiv:2008.11147 2020.
5. A. Meyer E. T. Barr C. Bird and T. Zimmermann "Today was a good day: The daily life of software developers" IEEE Transactions on Software Engineering 2019.
6. A. Felstead and G. Henseke "Assessing the growth of remote working and its consequences for effort well-being and work-life balance" New Technology Work and Employment vol. 32 no. 3 pp. 195-212 2017.
7. Naik GS, Gaur GS, Pal GK (2018) Effect of modified slow breathing exercise on perceived stress and basal cardiovascular parameters. Int J Yoga 11(1):53–58.
8. Meyer André N., Fritz Thomas, Murphy Gail C., and Zimmermann Thomas. 2014. Software developers' perceptions of productivity.
9. Johnson Brittany, Zimmermann Thomas, and Bird Christian. 2021. The effect of work environments on productivity and satisfaction of software engineers. IEEE Trans. Software Eng. 47, 4 (2021), 736–757.
10. Darja Smite: Changes in perceived productivity of software engineers during COVID-19 pandemic: The voice of evidence: Journal of Systems and Software: Volume 186, April 2022, 111197.